

What is Chlamydia?

Chlamydia is a STI that can infect both men and women. Most people who have chlamydia have no symptoms.

Symptoms in men may include:

- A burning sensation when urinating
- A discharge from the penis
- Painful or swollen testicles

Symptoms in women may include:

- Painful or burning sensation when urinating
- Abnormal vaginal discharge

People can become infected with Chlamydia during the following activities:

- Engaging in unprotected sex
- * This includes vaginal, anal, or oral sex

About 75% of women & 50% of men will not have symptoms of chlamydia.



1 in 5 people have a STI!

What is Syphilis?

Syphilis is a STI that can cause serious health problems if it is not treated.

People can become infected with Syphilis during the following activities:

- Direct contact with a syphilis sore during vaginal, anal, or oral sex
- Syphilis can spread from an infected mother to her unborn baby

It may not be obvious that a sex partner has syphilis as syphilis sores can be hidden in the vagina, anus, under the foreskin of the penis, or in the mouth. Unless you know that your partner has been tested and treated, you may be at risk of contracting syphilis.



Fort Riley Public Health
Building 7665 Normandy Drive
(785) 239-7323 / 7042



Public Health
Prevent. Promote. Protect.

★ U.S. ARMY Fort Riley, KS 

Walk-In Wellness

STI/STD Testing Program

IACH Lab M-F: 0730-1600

(Soldiers, Spouses, Beneficiaries 16 years & older)

Farrelly Lab M-F: 0730-1130

(Soldiers Only)

(Walk-in testing screens for the 5 diseases discussed in this handout; additional testing is available by PCM order. For more information, talk to your PCM or contact Fort Riley Public Health)

Walk-in 3-Site and Multiplex Vaginal Panel testing is available at Public Health on Fridays from 0800-1000

Book an Appointment!

(IACH Lab Only)



Type "walk-in STI check" in the PCM line

Ask about Expedited Partner Therapy!

What is HIV?

The Human Immunodeficiency Virus (HIV) can lead to Acquired Immunodeficiency Syndrome (AIDS) if not treated. HIV is a life-long condition.

People can become infected with HIV during the following activities:

- Unprotected sex
 - * This includes vaginal, anal, or oral sex
- Unsafe needle or syringe use
- Blood transfusions

Confidential counseling & education is provided by Army Public Health Nursing



Before having sex for the first time with a new partner, you and your partner should talk about your sexual history and consider getting tested for HIV and other STIs.

What is Gonorrhea?

Gonorrhea is a STI that can infect the genitals, rectum, and throat. Some men, and most women, with gonorrhea may have no symptoms.

Symptoms in men may include:

- A burning sensation when urinating
- A white, yellow, or green discharge from the penis
- Painful or swollen testicles (less common)

Symptoms in women may include:

- Painful or burning sensation when urinating
- Increased vaginal discharge

Drug-resistant strains of gonorrhea are increasing.

People can become infected with Chlamydia during the following activities:

- Engaging in unprotected sex
 - * This includes vaginal, anal, or oral sex



Get Peace of Mind!

What is Hepatitis C?

HEP C is a liver infection caused by a virus and can range from a mild illness lasting a few weeks to a serious, lifelong condition.

People can become infected with HEP C virus during the following activities:

- Sharing needles, syringes, or other equipment to prepare or inject drugs
- Child birth

Less commonly, a person can also get HEP C virus during the following activities:

- Needle stick injuries
- Sharing personal care items such as razors or toothbrushes
- Unprotected sex with an infected person
- Getting a tattoo or piercing in an unregulated setting

People with acute HEP C usually do not have symptoms or have mild symptoms. When symptoms do occur, they can include:

- Fever
- Loss of Appetite
- Fatigue
- Nausea
- Dark Urine
- Vomiting
- Abdominal Pain
- Joint Pain
- Jaundice